

Managing children who are sick, infectious, or with allergies

Policy statement

We aim to provide care for healthy children through preventing cross infection of viruses and bacterial infections and promote health through identifying allergies and preventing contact with the allergenic trigger.

Procedures for children who are sick or infectious

- If children appear unwell during the day – for example, if they have a temperature, sickness, diarrhoea or pains, particularly in the head or stomach – our manager call the parents and ask them to collect the child, or to send a known carer to collect the child on their behalf.
- If a child has a temperature, they are kept cool, by removing top clothing and sponging their heads with cool water, but kept away from draughts.
- The child's temperature is taken using a forehead thermometer strip, or an ear thermometer, kept in the first aid box.
- If the child's temperature does not go down and is worryingly high, then we may give them Calpol or another similar analgesic, after first obtaining verbal consent from the parent where possible. This is to reduce the risk of febrile convulsions, particularly for babies. Parents sign the medication record when they collect their child.
- In extreme cases of emergency, an ambulance is called and the parent informed.
- Parents are asked to take their child to the doctor before returning them to the setting; we can refuse admittance to children who have a temperature, sickness and diarrhoea or a contagious infection or disease.
- Where children have been prescribed antibiotics for an infectious illness or complaint, we ask parents to keep them at home for 48 hours before returning to the setting.
- After diarrhoea, we ask parents keep children home for 48 hours following the last episode.
- Some activities, such as sand and water play, and self-serve snacks where there is a risk of cross-contamination may be suspended for the duration of any outbreak.
- We a list of excludable diseases and current exclusion times.

Reporting of 'notifiable diseases'

- If a child or adult is diagnosed as suffering from a notifiable disease under the Health Protection (Notification) Regulations 2010, the GP will report this to Public Health England.

- When we become aware, or are formally informed of the notifiable disease, our manager informs Ofsted and contacts Public Health England, and act[s] on any advice given.

HIV/AIDS/Hepatitis procedure

HIV virus, like other viruses such as Hepatitis A, B and C, are spread through body fluids. Hygiene precautions for dealing with body fluids are the same for all children and adults. We :

- Wear single-use vinyl gloves and aprons when changing children's nappies, pants and clothing that are soiled with blood, urine, faeces or vomit.
- Bag soiled clothing for parents to take home for cleaning.
- Clear spills of blood, urine, faeces or vomit using mild disinfectant solution and mops; any cloths used are disposed of with the clinical waste.
- Clean any tables and other furniture, furnishings or toys affected by blood, urine, faeces or vomit using a disinfectant.

Nits and head lice

- Nits and head lice are not an excludable condition; although in exceptional cases we may ask a parent to keep the child away until the infestation has cleared.
- On identifying cases of head lice, we inform all parents ask them to treat their child and all the family if they are found to have head lice.
- If nits and head lice go untreated then the manager would discuss the situation with the parents/carers and seek professional advice on how to support the family, i.e. talk to their Health Visitor

Procedures for children with allergies

- When children start at the setting we ask their parents if their child suffers from any known allergies. This is recorded on the Registration Form.
- If a child has an allergy, we complete a risk assessment form to detail the following:
 - The allergen (i.e. the substance, material or living creature the child is allergic to such as nuts, eggs, bee stings, cats etc).
 - The nature of the allergic reactions (e.g. anaphylactic shock reaction, including rash, reddening of skin, swelling, breathing problems etc).
 - What to do in case of allergic reactions, any medication used and how it is to be used (e.g. Epipen).
 - Control measures - such as how the child can be prevented from contact with the allergen.
 - Review measures.
- This risk assessment form is kept in the child's personal file and a copy is displayed where our staff can see it.
- Generally, no nuts or nut products are used within the setting.
- Parents are made aware so that no nut or nut products are accidentally brought in, for example to a party.
- Life-saving medication and invasive treatments:

These include adrenaline injections (Epipens) for anaphylactic shock reactions (caused by allergies to nuts, eggs etc) or invasive treatments such as rectal administration of Diazepam (for epilepsy).

- We must have:
 - a letter from the child's GP/consultant stating the child's condition and what medication if any is to be administered;
 - written consent from the parent or guardian allowing our staff to administer medication; and
 - proof of training in the administration of such medication by the child's GP, a district nurse, children's nurse specialist or a community paediatric nurse.

- Key person for special needs children requiring assistance with tubes to help them with everyday living e.g. breathing apparatus, to take nourishment, colostomy bags etc.:
 - Prior written consent must be obtained from the child's parent or guardian to give treatment and/or medication prescribed by the child's GP.
 - They key person must have the relevant medical training/experience, which may include receiving appropriate instructions from parents or guardians.

Owner(S) name	Tarra Pitman and Carla Wilson
Owner(S) signature	
Date policy was rewritten	August 2018
Date policy is to be reviewed	August 2019